



## THE PIONEERING CONTRIBUTIONS AND LEGACY OF WALDO H. ZAGWIJN (1928–2018)

Jean-Pierre Suc<sup>1</sup>, Speranta-Maria Popescu<sup>2</sup>, Adele Bertini<sup>3</sup>,  
Martin J. Head<sup>4</sup>, Philip L. Gibbard<sup>5</sup>, Filomena Diniz<sup>6</sup>

<sup>1</sup> Sorbonne Université, CNRS-INSU, Institut des Sciences de la Terre, Paris, France

<sup>2</sup> GeoBioStratData, Consulting, Rillieux la Pape, France

<sup>3</sup> Dipartimento di Scienze della Terra, Università di Firenze, Firenze, Italy

<sup>4</sup> Department of Earth Sciences, Brock University, St. Catharines, Ontario, Canada

<sup>5</sup> Scott Polar Research Institute, University of Cambridge, Cambridge, UK

<sup>6</sup> Departamento de Geologia, Universidade de Lisboa, Lisboa, Portugal

Corresponding author: J.-P. Suc <[jeanpierre.suc@gmail.com](mailto:jeanpierre.suc@gmail.com)>

**ABSTRACT:** Professor Dr. Waldo H. Zagwijn passed away on 26 June, 2018, in his ninetieth year. His thesis research (Zagwijn, 1960) stimulated palynological investigations into the Pliocene and early Quaternary, and caused a fundamental shift in our understanding of the transition from the warm and relatively stable climate of the late Neogene to the extreme glacial–interglacial oscillations of the early Quaternary. Distinct in several aspects from the usual methodologies of that time, Zagwijn’s approach can be summarized as:

- focusing on establishing the botanical identification of pollen grains, following the guiding approach of his supervisor F. Florschütz, using botanical nomenclature to highlight these identifications and clarify relationships between botany and the earlier artificial pollen nomenclature;
- comprehensive and persistent attention to plant macroremains (fruits, seeds, etc.; Zagwijn, 1990) as a mean of integrating classical palaeobotany with modern palynology;
- palaeoenvironmental interpretations informed by a deep understanding of modern plant ecology and associations;
- leading investigations in the field, and paying special attention to stratigraphic correlations between (onshore and offshore) boreholes and exposed sections in quarries, using a range of available proxies including heavy mineral analysis.

Waldo H. Zagwijn was a visionary in realizing how modern palynology can be used for vegetational and climatic reconstructions in the Neogene, and indeed as far back as Paleogene times.

**Keywords:** pollen analysis, palaeovegetation, climatostratigraphy, Pliocene–Pleistocene boundary, glacials–interglacials

### 1. MODERN PALYNOLOGY AND ITS CONTRIBUTION TO GLOBAL CLIMATOSTRATIGRAPHY

Using the methodological novelties detailed above, Zagwijn’s thesis identified a pronounced cooling event (marked by the establishment of tundra-like vegetation in The Netherlands) separating the Reuverian and Tiglian continental regional stages, and he identified this as the Praetiglian, a stage named earlier by Van der Vlerk & Florschütz (1953) from the classic Dutch-German border area. Zagwijn had in fact discovered the onset of pronounced glacial–interglacial cycles in the Northern Hemisphere. This immediately brought his work to the attention of the international scientific community. This first major contribution of Zagwijn was to show how the evolution of vegetation and climate of North Europe progressed cyclically from subtropical to warm-temperate and glacial conditions. It resulted in the establishment of a robust climatostratigraphy with the possibilities of long-distance correlation (Van der Hammen et al., 1971), based on botanical pollen classification and counting, rather than the now outdated palynostratigraphic method constructed on the presence or absence of fictitious species.

Aiming to obtain independent chronological control

for the record of climate changes, Zagwijn proposed correlations with the Mediterranean succession, and this included locating the Pliocene–Pleistocene boundary, which had been under discussion since 1948, at the Piacenzian–Calabrian contact (Zagwijn, 1974, 1975a). Simultaneously, Zagwijn explored magnetostratigraphy as a mean to calibrate climate evolution with the global timescale (Zagwijn, 1975b) and paid special attention to mammal biochronology in support of regional correlations (e.g., Van der Meulen & Zagwijn, 1974). Oxygen isotope curves showed that the Praetiglian glacial stage occurred in what was then considered the late Pliocene (Shackleton & Opdyke, 1977; Blanc et al., 1983; Shackleton et al., 1984), as supported by subsequent pollen records in the northwestern Mediterranean region (Cravatte & Suc, 1981; Suc & Cravatte, 1982; Suc, 1984). As a consequence, a reliable climatostratigraphic relationship between The Netherlands and the Mediterranean was established and the first significant cooling in the Northern Hemisphere (onset of tundra-like vegetation to the north, and steppe vegetation to the south) was placed near the Gauss–Matuyama reversal (Suc & Zagwijn, 1983). The discovery confirmed the value of pollen records for long-distance correlation and strengthened the chronological framework for climate change in




Fig. 1 - Example of climatostratigraphic correlations at the European scale between six long pollen records from the earliest Pliocene to earliest Pleistocene and chronologic calibration of the Northwestern European climatic succession (modified from Popescu et al., 2010). Pollen groups (thermic classification: Nix, 1982): 1, Megathermic elements (*Canthium*, Sapindaceae, *Bombax*, etc.); 2, Mega-mesothermic elements (*Taxodium*-type, *Sequoia*-type, Arecaceae, *Engelhardia*, *Distylium*, *Microtropis fallax*, etc.); 3, *Cathaya* (= *Pinus* haploxylon-type at Wólka Ligezawska); 4, Mesothermic elements (*Quercus*, *Carya*, *Pterocarya*, *Liquidambar*, *Carpinus*, *Ulmus*, *Zelkova*, etc.); 5, *Pinus*; 6, Meso-microthermic trees (*Cedrus*, *Tsuga*, *Keteleeria*); 7, Microthermic trees (*Abies*, *Picea*); 8, Elements of no significance (Ranunculaceae, Rosaceae, etc.); 9, Cupressaceae (*Cupressus*-*Juniperus*-type); 10, Mediterranean xerophytes (*Olea*, *Pistacia*, *Ceratonia*, *Quercus ilex*-type, *Rhamnus*, *Cistus*, etc.); 11, Herbs (Amaranthaceae, Asteraceae, Poaceae, Apiaceae, *Rumex*, Boraginaceae, *Convolvulus*, Cyperaceae, Caryophyllaceae, etc.); 12, Steppe elements (*Artemisia*, *Ephedra*, *Hippophae*); 13, Ericaceae.

Europe (Zagwijn, 1986; Gibbard et al., 1991; Suc et al., 1995; Pontini & Bertini, 2000; Bertini, 2001; Popescu et al., 2010) (Fig. 1). In addition his work provided a firm foundation for reconstructing the evolution of the North Sea region throughout the late Cenozoic.

## 2. THE LOWERING OF THE PLIOCENE-PLEISTOCENE BOUNDARY FROM 1.8 TO 2.58 MA

The age of Praetiglian cooling having been established within the late Pliocene (at that time, the Pliocene extended to 1.8 Ma), the International Commission of Stratigraphy (ICS) approved the creation of a third stage for the Pliocene, the Gelasian (2.6–1.8 Ma), sandwiched between the Piacenzian Stage (3.6–2.6 Ma) and the intact Calabrian Stage (starting at 1.8 Ma) (Rio et al., 1998), aiming to avoid the prospect of lowering the Pliocene–Pleistocene boundary from 1.8 to 2.6 Ma (Suc et al., 1997). However, the Quaternary community consolidated its position and, despite a disappointing setback in 1997 (for full details, see: Head & Gibbard, 2015), succeeded on 29 June 2009 in obtaining formal ratification from the executive committee of the IUGS to lower the Pliocene–Pleistocene boundary to the base of the Gelasian Stage with an age of 2.58 Ma (Gibbard & Head, 2010; Gibbard et al., 2010) (Fig. 2).

Waldo H. Zagwijn was a strong and vocal advocate of a Pliocene–Pleistocene (and Neogene–Quaternary) boundary that coincided with the base of the Praetiglian, and he readily offered arguments in support of this view, whether to opponents or supporters. Zagwijn was also convinced that Quaternary time

### Pre-2009 time scale

SYSTEM PERIOD	SERIES EPOCH	STAGE AGE
QUATERNARY	HOLOCENE	
	PLEISTOCENE	Unnamed
		Unnamed
NEOGENE	PLIOCENE	Calabrian
		Gelasian
	Piacenzian	
	Zanclean	

### Current time scale

SYSTEM PERIOD	SERIES EPOCH	STAGE AGE	GSSP
QUATERNARY	HOLOCENE	Meghalayan	0.0042
		Northgrippian	0.0082
		Greenlandian	0.0117
	PLEISTOCENE	Unnamed	0.129
		Unnamed	0.773
		Calabrian	1.80
		Gelasian	2.58
NEOGENE	PLIOCENE	Piacenzian	3.60
		Zanclean	5.33

Fig. 2 - The geological time scale as it existed before the base of the Pleistocene was lowered to that of the Gelasian in 2009, and the current IUGS/ICS geological time scale. [IUGS/ICS geological time scale.]

was distinct and that it should not be incorporated into the Neogene (Fig. 2), as some of his countrymen had advocated for several decades. There can be no question that, at the age of 81, he felt this IUGS decision to




Fig. 3 - Climatic curves resulting from pollen records compared to oxygen isotope curves. A) The Netherlands climatic curve matched to an oxygen isotope curve (modified from Zagwijn, 1989); B) Focus on the latest Praetiglian to early Waalian pollen ratio 'thermophilous/steppe elements' from the Mediterranean region (Croton, Calabrian Stage) compared with the LR04 benthic  $\delta^{18}\text{O}$  stack (modified from Suc et al., 2010, 2018).

be the ultimate recognition of his work. This was not only a vindication of his concept but an acknowledgment that climate (and of course palynology) should play a prominent role in formal stratigraphy, at least for the Quaternary.

### 3. EXTENDING THE NUMBER OF GLACIAL-INTERGLACIAL CYCLES, AND DRAWING PALYNOLOGY CLOSER TO ISOTOPE STRATIGRAPHY

After defining how respectively to characterise glacial and interglacials by pollen assemblages and then delimit them (Zagwijn, 1957), Zagwijn (1960) proposed a climatic subdivision of the Pleistocene of The Netherlands that has been continuously refined to this day by the discovery of new phases (Zagwijn, 1989, 1998; Fig. 2A) extendable over the entire southern North Basin (Gibbard et al., 1991). This evolution is explained by the discontinuous character of sedimentation in the delta to flood-plain environmental contexts in which Zagwijn worked. Integration was achieved by piecing together sequences based on the integration of multidisciplinary litho- and biostratigraphical investigations from across the country (Zagwijn & Doppert, 1978; Zagwijn, 1989). Keen to compare his pollen data with

oxygen isotope records, Zagwijn (1985) constructed a curve of estimated summer (July) mean temperatures in The Netherlands from pollen-analytical data, and data on cryoturbatic structures (Fig. 2A; for details, see: Zagwijn, 1985). Zagwijn's curve was validated as representative of climatic megacycles by Kukla & Cilek (1996).

Condensing pollen data into a single curve is a crucial challenge because the progressive enrichment of pollen floras by many new taxa over time complicates interpretation. In the Mediterranean region, where temperature and precipitation critically combine to characterise climate change, a pollen index is displayed by the ratio of 'thermophilous plants/steppe elements', which provides a curve directly comparable to oxygen isotope records (Fig. 3B; Suc et al., 2010). The curves illustrate the great variability of climate during each phase of a megacycle as shown for the Tiglian and Eburonian stages (Fig. 3B). For example, the Praetiglian as characterised by W.H. Zagwijn with few or no minor subdivisions, has progressively been shown to abound in secondary climatic fluctuations (Suc, 1984; Pontini & Bertini, 2000; Bertini, 2001; Donders et al., 2018) and may have a complex lateral expression owing to local palaeoenvironmental variation (Donders et al., 2007). Following the path well trodden in The Netherlands, sustained effort



Fig. 4 - Waldo H. Zagwijn (to the right) receiving the Penck Medal in 1973, photographed by the organizer.

has been directed at collecting new pollen data and synthesizing with existing records in the Mediterranean region to enhance our understanding of vegetational and climatic evolution (Suc & Popescu, 2005; Leroy, 2007; Bertini, 2010; Popescu et al., 2010; Combourieu-Nebout et al., 2015; Suc et al., 2018). Finally, pollen transfer functions are now being used to reconstruct past climate from locations across Europe (Fauquette et al., 2007).

#### 4. AN IMPRESSIVE PERSON AND A BRILLIANT CAREER

Anyone who has known Waldo Zagwijn remembers his humility and constant attention to others. His intelligence shone through his sparkling eyes. Always affable and imbued with a keen sense of detail, Zagwijn was immensely interested in training young geologists and palynologists as illustrated by his commitment as a Professor at the Free University Amsterdam.

Well into retirement, Waldo H. Zagwijn remained keenly interested in biology and geology, traits that guided his eminent career at the Geological Survey of The Netherlands. Crowned with numerous distinctions including Life Fellowship of INQUA, royal appointment as an officer of the Order of Orange-Nassau, award and recipient of the Penck Medal (Fig. 4), he was elected in 1980 as a member of the Royal Academy of Sciences of

The Netherlands (*Koninklijke Nederlandse Akademie van Wetenschappen*). On the occasion of his retirement, the Accademy celebrated his life's work on October 1993 with an international symposium and a volume including his complete bibliography and many dedicated papers (Hengreen & van der Valk, 1995).

#### REFERENCES

- Bertini A. (2001) - Pliocene climatic cycles and altitudinal forest development from 2.7 Ma in the Northern Apennines (Italy): evidence from the pollen record of the Stirone section (5.1 to 2.2 Ma). *Geobios*, 34 (3), 253-265.
- Bertini A. (2010) - Pliocene to Pleistocene palynoflora and vegetation in Italy: State of the art. *Quaternary International*, 225, 5-24.
- Blanc P.L., Fontugne M.R., Duplessy J.C. (1983) - The time-transgressive initiation of boreal ice-caps: continental and oceanic evidence reconciled. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 42, 211-224.
- Combourieu-Nebout N., Bertini A., Russo Ermolli E., Peyron O., Klotz S., Montade V., Fauquette S., Allen J., Fusco F., Goring S., Huntley B., Joannin S., Lebreton V., Magri D., Martinetto E., Orain R., Sadori L. (2015) - Climate changes in the central Mediterranean and Italian vegetation dynamics

- since the Pliocene. Review of Palaeobotany and Palynology, 218, 127-147.
- Cravatte J., Suc J.-P. (1981) - Climatic evolution of North-Western Mediterranean area during Pliocene and Early Pleistocene by pollen-analysis and forams of drill Autan 1. Chronostratigraphic correlations. Pollen et Spores, 23(2), 247-258.
- Diniz F. (1984) - Etude palynologique du bassin pliocène de Rio Maior. Paléobiologie Continentale, 14, 259-267.
- Donders T.H., Kloosterboer van Hoeve M.L., Westerhoff W., Verreussel R.H.C.M., Lotter A.F. (2007) - Late Neogene continental stages in NW Europe revisited. Earth Science Reviews, 85, 161-186.
- Donders T.H., van Helmond N.A.G.M., Verreussel R., Munsterman D., ten Veen J., Speijer R.P., Weijers J.W.H., Sangiorgi F., Peterse F., Reichert G.J., Sinninghe Damsté, J.S., Lourens, L., Kuhlmann, G., Brinkhuis, H. (2018) - Land-sea coupling of early Pleistocene glacial cycles in the southern North Sea exhibit dominant Northern Hemisphere forcing. Climate of the Past, 14, 397-411.
- Fauquette S., Suc J.-P., Jiménez Moreno G., Micheels A., Jost A., Favre E., Bachiri Taoufiq N., Bertini A., Clet Pellerin M., Diniz F., Farjanel G., Feddi N., Zheng, Z. (2007) - Latitudinal climatic gradients in Western European and Mediterranean regions from the Mid-Miocene (c. 15 Ma) to the Mid-Pliocene (c. 3.5 Ma) as quantified from pollen data. In "Deep-Time Perspectives on Climate Change. Marrying the Signal from Computer Models and Biological Proxies", Williams M., Haywood A., Gregory J., Schmidt D.N. (Eds.). The Micropaleontological Society, The Geological Society, London, Special Publications, 481-502.
- Gibbard P.L., Head M.J. (2010) - The newly-ratified definition of the Quaternary System/Period and redefinition of the Pleistocene Series/Epoch, and comparison of proposals advanced prior to formal ratification. Episodes, 33, 152-158.
- Gibbard P.L., Head M.J., Walker M.J.C., and the Subcommission on Quaternary Stratigraphy (2010) - Formal ratification of the Quaternary System/Period and the Pleistocene Series/Epoch with a base at 2.58 Ma. Journal of Quaternary Science, 25(2), 96-102.
- Gibbard P.L., West R.G., Zagwijn W.H., Balson P.S., Burger A.W., Funnell B.M., Jeffery D.H., de Jong J., van Kolfschoten T., Lister A.M., Meijer T., Norton P.E.P., Preece R.C., Rose J., Stuart A.J., Whiteman C.A., Zalasiewicz J.A. (1991) - Early and early middle Pleistocene correlations in the southern North Basin. Quaternary Science Reviews, 10, 23-52.
- Head M.J., Gibbard P.L. (2015) - Formal subdivision of the Quaternary System/Period: Past, present, and future. Quaternary International, 383, 4-35.
- Herngreen G.F.W., van der Valk L. (1995) (Eds.) - Neogene and Quaternary geology of North-West Europe. Contributions on the occasion of Waldo H. Zagwijn's retirement. Mededelingen Rijks Geologische Dienst, 52, 510 p.
- Kukla G., Cílek V. (1996) - Plio-Pleistocene megacycles: record of climate and tectonics. Palaeogeography, Palaeoclimatology, Palaeoecology, 120, 171-194.
- Leroy S.A.G. (2007) - Progress in palynology of the Gelasian-Calabrian Stages in Europe; Ten messages. Revue de Micropaléontology, 50, 293-308.
- Lisiecki L.E., Raymo M.E. (2005) - A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. Paleoceanography, 20, PA1003.
- Nix H. (1982) - Environmental determinants of biogeography and evolution in Terra Australis, In "Evolution of the Flora and fauna of Arid Australia", Barker W.R. & Greenslade P.J.M. (Eds.), Peacock Publishing, Frewville, 47-66.
- Pontini R., Bertini A. (2000) - Late Pliocene vegetation and climate in Central Italy: high-resolution pollen analyses from the Fosso Bianco succession (Tiberino Basin). Geobios, 33(5), 519-526.
- Popescu S.M. (2001) - Repetitive changes in Early Pliocene vegetation revealed by high-resolution pollen analysis: revised cyclostratigraphy of southwestern Romania. Review of Palaeobotany and Palynology, 120(3-4), 181-202.
- Popescu S.M., Biletekin D., Winter H., Suc J.-P., Melinte-Dobrinescu M.C., Klotz S., Combourieu Nebout N., Rabineau M., Clauzon G., Deaconu F. (2010) - Pliocene and Lower Pleistocene vegetation and climate changes at the European scale: Long pollen records and climatostratigraphy. Quaternary International, 219, 152-167.
- Rio D., Sprovieri R., Castradori D., Di Stefano E. (1998) - The Gelasian Stage (Upper Pliocene): A new unit of the global standard chronostratigraphic scale. Episodes, 21, 82-87.
- Shackleton N.J., Backman J., Zimmerman H., Kent D.V., Hall M.A., Roberts D.G., Schnitker D., Baldauf J.G., Desprairies A., Homrighausen R., Huddleston P., Keene J.B., Kaltenback A.J., Krumsiek K.A.O., Morton A.C., Murray J.W., Westberg Smith J. (1984) - Oxygen isotope calibration of the onset of ice-rafting and history of glaciation in the North Atlantic region. Nature, 307, 620-623.
- Shackleton N.J., Opdyke N.D. (1976) - Oxygen-isotope and paleomagnetic stratigraphy of Pacific Core V28-239 Late Pliocene to Latest Pleistocene. Geological Society of America Memory, 145, 449-464.
- Shackleton N.J., Opdyke N.D. (1977) - Oxygen isotope and palaeomagnetic evidence for early Northern Hemisphere glaciation. Nature, 270, 216-219.
- Suc J.-P. (1984) - Origin and evolution of the Mediterranean vegetation and climate in Europe. Nature, 307, 429-432.
- Suc J.-P., Bertini A., Leroy S.A.G., Suballyova D. (1997) - Towards the lowering of the Pliocene/Pleistocene boundary to the Gauss-Matuyama reversal. Quaternary International, 40, 37-42.
- Suc J.-P., Combourieu-Nebout N., Seret G., Popescu S.-M., Klotz S., Gautier F., Clauzon G., Westgate J., Sandhu A.S. (2010) - The Crotone series: a synthesis and new data. Quaternary International, 219, 121-133.
- Suc J.-P., Cravatte J. (1982) - Etude palynologique du

- Pliocène de Catalogne (nord-est de l'Espagne). *Paléobiologie Continentale*, 13(1), 1-31.
- Suc J.-P., Diniz F., Leroy S., Poumot C., Bertini A., Dupont L., Clet M., Bessais E., Zheng Z., Fauquette S., Ferrier J. (1995) - Zanclean (~ Brunssumian) to early Piacenzian (~ early-middle Reuverian) climate from 4° to 54° north latitude (West Africa, West Europe and West Mediterranean areas). *Mededelingen Rijks Geologische Dienst*, 52, 43-56.
- Suc J.-P., Popescu S.-M. (2005) - Pollen records and climatic cycles in the North Mediterranean region since 2.7 Ma. In "Early-Middle Pleistocene Transitions: The Land-Ocean Evidence", Head M.J. & Gibbard P.L. (Eds.), Geological Society of London, Special Publication, 247, 147-158.
- Suc J.-P., Popescu S.-M., Fauquette S., Bessedik M., Jiménez Moreno G., Bachiri Taoufiq N., Zheng Z., Médail F. (2018) - Reconstruction of Mediterranean flora, vegetation and climate for the last 23 million years based on an extensive pollen dataset. *Ecologia Mediterranea*, 44, 2, in press.
- Suc J.-P., Zagwijn W.H. (1983) - Plio-Pleistocene correlations between the northwestern Mediterranean region and northwestern Europe according to recent biostratigraphic and palaeoclimatic data. *Boreas*, 12, 153-166.
- Van der Hammen T., Wijmstra T.A., Zagwijn W.H. (1971) - The floral record of the late Cenozoic of Europe. In "The late Cenozoic glacial ages", Turekian K.K. (Ed.), Yale University press, New Haven and London, 391-424.
- Van der Meulen A.J., Zagwijn W.H. (1974) - *Microtus (Allophaiomys) pliocaenicus* from the Lower Pleistocene near Brielle, The Netherlands. *Scripta Geologica*, 21, 1-12.
- Van der Vlerk I.M., Florschütz F. (1953) - The palaeontological base of the subdivision of the Pleistocene in the Netherlands, *Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen, Afdeling Natuurkunde*, 1e Reeks XX, 2, 1-58.
- Zagwijn W.H. (1957) - Vegetation, climate and time-correlations in the Early Pleistocene of Europe. *Geologie en Mijnbouw, (N.S.)*, 19, 233.
- Zagwijn W.H. (1960) - Aspects of the Pliocene and early Pleistocene vegetation in The Netherlands. *Mededelingen van de Geologische Stichting, (C)*, 3(1), 5, 4-78.
- Zagwijn W.H. (1974) - The Plio-Pleistocene boundary in western and southern Europe. *Boreas*, 3, 75-97.
- Zagwijn W.H. (1975a) - A model-theory for the Pliocene/Pleistocene boundary determination, based on past climatic changes. In "Late Neogene Epoch boundaries", Saito T. & Burckle L.H. (Eds.), *Micro-paleontology, Special Publication*, 71-74.
- Zagwijn W.H. (1975b) - Variations in climate as shown by pollen analysis, especially in the Lower Pleistocene of Europe. In "Ice ages: ancient and modern", Wright A.E., Moseley F. (Eds.), 137-152.
- Zagwijn W.H. (1985) - An outline of the Quaternary stratigraphy of the Netherlands. *Geologie en Mijnbouw*, 64, 17-24.
- Zagwijn W.H. (1986) - Plio-Pleistocene climatic change: evidence from pollen assemblages. *Memorie della Società Geologica Italiana*, 31, 145-152.
- Zagwijn W.H. (1989) - The Netherlands during the Tertiary and the Quaternary: A case history of Coastal Lowland evolution. *Geologie en Mijnbouw*, 68, 107-120.
- Zagwijn W.H. (1990) - Subtropical relicts in the Pliocene flora of Brunssum (The Netherlands). *Geologie en Mijnbouw*, 69, 219-225.
- Zagwijn W.H., Doppert J.W.Chr. (1978) - Upper Cenozoic of the southern North Sea Basin: Palaeoclimatic and palaeogeographic evolution. *Geologie en Mijnbouw*, 57(4), 577-588.
- Zagwijn W.H. (1998) - Borders and boundaries: a century of stratigraphical research in the Tegelen - Reuver area of Limburg (The Netherlands). In "The Dawn of the Quaternary", Van Kolfschoten Th., Gibbard P.L. (Eds.), *Proceedings of the SEQS-EuroMan Symposium 1996*, 60. *Mededelingen Nederlands Instituut voor Toegepaste geowetenschappen TNO*, 19-34.

Ms. received: November 1, 2018

Final text received: November 27, 2018